

B'SYS GmbH

CHO K_v10.1 Cell Line

Cell culture conditions

© B'SYS GmbH

TABLE OF CONTENTS

1.	PRODUCT SHIPMENT	3
1.1.	PRODUCT FORMAT	3
1.2.	MYCOPLASMA CERTIFICATE.....	3
2.	CELL CULTURE CONDITIONS	4
2.1.	GENERAL	4
2.2.	RECOMMENDED COMPLETE MEDIUM	4
2.3.	ANTIBIOTICS	4
2.4.	THAWING CELLS	4
2.5.	SPLITTING CELLS.....	5
2.6.	FREEZING MEDIUM	5
2.7.	FREEZING CELLS	5
2.8.	STABILITY OF CHO Kv10.1 CELLS	5
3.	Kv10.1 SEQUENCE	5
3.1.	ACCESSION NUMBER NP_002229.1	5
4.	CONTACT INFORMATION.....	6
4.1.	CONTACT ADDRESS FOR TECHNICAL SUPPORT AND ORDERING INFORMATION	6

1. PRODUCT SHIPMENT

1.1. Product Format

CHO cells stably transfected with recombinant Kv10.1 potassium channel:

- 0.75 mL aliquots of frozen cells at 1 E+06 cells/mL
- Cells are frozen in complete medium with 10% DMSO
- Cells are frozen at passage number, see vial

1.2. Mycoplasma Certificate

B'SYS periodically tests cells for presence of mycoplasma by means of highly sensitive PCR based assays. All delivered cells are free of mycoplasma.

2. CELL CULTURE CONDITIONS

2.1. General

CHO Kv10.1 cells are incubated at 37°C in a humidified atmosphere with 5% CO₂ (rel. humidity > 95%). The cells are continuously maintained and passaged in sterile culture flasks containing F12 (HAM) medium supplemented with 10% foetal bovine serum, 1.0% Penicillin/Streptomycin solution and 300 µg/mL Hygromycin. The CHO Kv10.1 cells are passaged at a confluence of about 80%.

- All solutions and equipment coming in contact with the cells must be sterile.
- Use proper sterile technique and work in a laminar flow hood.
- Be sure to have frozen cell stocks at hand before starting experiments.
- Cells should be split every 2-3 days at 70% - 80% confluence at 1:3 to 1:5 ratio.

Table 1: Cell culture reagents

Product	Supplier	Order number
Nutrient mixture F-12 Ham (F12 Ham)	Sigma-Aldrich	N6658
Fetal Bovine Serum (FBS)	Gibco	10270-106
Penicillin / Streptomycin (100x)	Gibco	10378-016
Phosphate Buffered Saline (PBS, without Ca ²⁺ and Mg ²⁺)	Sigma-Aldrich	D8537
Hygromycin B (50 mg/mL)	Gibco	10687010
Detachin	Genlantis	T100T100
Trypsin EDTA (10x)	Sigma-Aldrich	T4174
DMSO	Sigma-Aldrich	D2438

For the preparation of 1X Trypsin/EDTA, the 10X solution is diluted in PBS (without Ca²⁺ and Mg²⁺), aliquoted and stored in the freezer.

2.2. Recommended Complete Medium

- 500 mL F12 (HAM) with L-Glutamine
- 10% FBS
- 1.0% Penicillin/Streptomycin

2.3. Antibiotics

- To cultivate CHO Kv10.1 cells, an antibiotic pressure of 300 µg/mL must be used.

Remark: The permanent application of high antibiotic pressure has no effect on current density.

2.4. Thawing Cells

- Remove vial of cells from liquid nitrogen and thaw quickly at 37°C.
- Decontaminate outside of vial with 70% Ethanol.
- Transfer cells to a T-25 culture flask containing 5 mL complete medium
- Incubate cells at 37°C for at least 4-6 hours to allow the cells to attach to the bottom of the flask
- Once cells attach to the bottom of the flask and look healthy, aspirate off the medium and replace with 5 mL complete medium containing selection antibiotics for cultivation (see 2.3)
- To check whether cells are attached properly, the flask can be gently moved while looking under the microscope
- If 48h after thawing the confluency is below 50%, replace the medium in the flask with fresh medium containing antibiotics

- Incubate cells at 37°C and check them daily until 50% to 80% confluency is reached.

2.5. Splitting Cells

- When cells are 50% to 80% confluent remove complete medium.
- Wash cells with 1xPBS to remove excess medium
- Add 0.5 mL to 1 mL Detachin (or 1x Trypsin/EDTA) and incubate for 2 min at 37°C.
- Detach cells by gently tapping the sides of the flask add complete medium and pipet up and down to break clumps of cells.
- Passage cells into new flask with complete medium and antibiotics at 1:3 to 1:5 ratio.
- Use remaining suspension for counting the cells.

2.6. Freezing Medium

- Mix 0.9 mL fresh complete medium and 0.1 mL DMSO for every 1 mL freezing medium.
- Sterilize freezing medium by means of appropriate micro filter (0.1 µm – 0.2 µm).

2.7. Freezing Cells

- Prepare fresh freezing medium and keep it on ice.
- Cells should have 80% - 90% confluency prior to freezing.
- Remove the complete medium and wash cells with 1xPBS.
- Add 0.5 mL to 1 mL Detachin (or 1x Trypsin/EDTA) and incubate for 2 min at 37°C
- Detach cells by gently tapping the sides of the flask, add complete medium and pipet up and down to break clumps of cells.
- Pellet cells at 200 g using a centrifuge and carefully aspirate off medium.
- Resuspend cells at a density of approximately 1.0 E+06 cells per mL with fresh freezing medium.
- Aliquot 0.75 mL of cell suspension into each cryovial.
- Overnight incubate cells in a polystyrene box at –80°C.
- The next morning transfer cryovial in liquid nitrogen tank for long-term storage.

2.8. Stability of CHO Kv10.1 cells

CHO Kv10.1 cells stably express functionally active Kv10.1 potassium channels over 10 passages. Under recommended cell culture conditions, no variation in current density was observed during this time.

3. Kv10.1 SEQUENCE

3.1. Accession Number NP_002229.1

Cloned cDNA is codon optimized and encodes for the protein of the human potassium voltage-gated channel subfamily H member 1, isoform 2

4. CONTACT INFORMATION

4.1. Contact Address for Technical Support and Ordering Information

- B'SYS GmbH
Technology Center Witterswil
Benkenstrasse 254
4108 Witterswil
Switzerland

Tel: +41 61 721 77 44

Fax: +41 61 721 77 41

Email: info@bsys.ch

Web: www.bsys.ch